

Elevated skin tissue cholesterol levels and myocardial infarction

Dennis L. Sprecher*, Gregory L. Pearce

University of Pennsylvania, Department of Experimental Therapeutics, Desk C51, Philadelphia, PA 19104, USA

Received 29 April 2004; received in revised form 12 October 2004; accepted 3 January 2005

Available online 8 March 2005

Abstract

Background: Skin cholesterol has been associated with coronary artery disease, extent of angiographic disease and inflammatory markers such as hs-CRP. Based on these findings we sought to determine whether skin cholesterol was associated with myocardial infarction (MI).

Methods: Patients ($N = 649$) underwent diagnostic catheterization and concurrent skin cholesterol measurement. History of MI was determined at the time of hospitalization.

Results: Patients with a history of MI ($n = 225$, 35%) had significantly higher skin cholesterol than those without MI (127 ± 29 versus 120 ± 20 , $p = 0.002$). The odds ratio for high skin cholesterol (for MI) was 1.6 (95% CI = 1.1, 2.6; $p = 0.01$) after adjustment for traditional risk and extent of angiographic disease.

Conclusion: Skin cholesterol may indicate increased risk of coronary-related events rather than simply the presence of angiographic narrowing. © 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Coronary artery disease; Skin; Myocardial infarction; Angiography

1. Introduction

Skin cholesterol, a surrogate for tissue cholesterol, has long been associated with coronary artery disease (CAD) [1] and more recently with the degree of angiographic stenosis [2]. Further, skin cholesterol has been found to be associated with inflammatory markers (VCAM and CRP) [3]. We would anticipate therefore that skin cholesterol may be associated with acute events, beyond any association with coronary arterial narrowing.

2. Methods

Six hundred and forty-nine patients, not on lipid lowering medications, undergoing non-emergency diagnostic catheterization had SkinT_c measured and baseline risk data recorded. The patient population was recruited consecutively from three separate sites, two in Canada (St. Michaels Hospital (SMH), $n = 323$; Trillium Health Center (THC), $n = 179$) and

one in the United States (Cleveland Clinic Foundation (CCF), $n = 147$). Technicians at each site were trained in an identical manner to measure SkinT_c. All catheterization interpretations were performed manually and independently at each site. The major indications for the catheterization was anginal symptoms (>90%) and positive stress test (>60%, >100% because some had both). Institutional Review Boards approved the study protocols at each site and all subjects provided informed consent prior to study enrollment.

Within 24 h prior to the angiographic procedure, SkinT_c measurement was performed. The non-biopsy based skin-cholesterol assay system (cholesterol 1, 2, 3TM), which uses a synthetic copolymer conjugated with digitonin and horseradish peroxidase to bind and quantify SkinT_c has been recently described [4]. Cholesterol is assayed only within the stratum corneum. Repetitive readings provide a mean day-to-day coefficient of variation of 3–8% with a between hand average reproducibility of 95%. The entire process requires about 3–4 min.

Extent of disease was defined as the number of vessels with $\geq 50\%$ stenosis and patients were further classified as having angiographic disease if the extent of disease was at least “one”. Traditional CAD risk was characterized by the

* Corresponding author. Tel.: +1 215 662 9993; fax: +1 215 243 3255.

E-mail address: dennis.sprecher@uphs.upenn.edu (D.L. Sprecher).




Fig. 1. Mean SkinT_c (with standard error) by MI status, CABG status and MI or CABG status.

Framingham global risk score as reported by Wilson et al. [5]. Patients without and with a history of myocardial infarction (MI) (recorded from patient history and chart review) were compared with chi-square tests for categorical measures and unpaired *t*-tests for continuous measures. High SkinT_c was defined as the upper quartile of the distribution (>143). Logistic regression techniques were used to estimate the relative risk of MI and (since coronary artery bypass surgery (CABG) subjects are enriched with those experiencing recent progressive angina), the combination of MI and/or CABG with odds ratios (OR), 95% confidence intervals (CI) and *p*-values.

3. Results

Among 649 patients scheduled for angiography, 225 (35%) had a history of MI, 50 (8%) had a history of coronary artery bypass surgery and 240 (37%) had a history of either MI or CABG. Presentation characteristics for patients without and with history of MI are found in Table 1. SkinT_c was significantly higher for patients with history of MI, CABG or either (Fig. 1).

High SkinT_c was associated with an increased rate of prior MI (OR = 1.5, CI = 1.1–2.9, *p* = 0.03). This association remained clear after adjustment Framingham risk and extent of disease (OR = 1.5, CI = 1.0–2.2, *p* = 0.05). The un-

adjusted OR was 1.7 (CI = 1.2–2.5, *p* = 0.003) for those patients with a history of MI and/or CABG. After adjustment for extent of disease and traditional risk burden the OR for high SkinT_c remained statistically significant (OR = 1.8, CI = 1.2–2.6, *p* = 0.005).

4. Discussion

In our study population of subjects scheduled for angiography, patients with a history of MI had significantly higher skin cholesterol levels. Patients with SkinT_c in the highest quartile were 1.5 times as likely to have had an MI and nearly twice as likely to have had an MI or CABG. These associations remain essentially unaffected by adjustment for extent of disease and traditional risk burden.

Based on the reported modest association between SkinT_c and extent of angiographic disease [2], as well as the association with markers of inflammation (e.g., VCAM [3]), we anticipated a relation between SkinT_c and acute events, outside of the association with extent of disease alone. This would be in line with the known connection between inflammatory markers and acute coronary events [6]. We now know that SkinT_c was correlated with history of MI, even with adjustment for level of angiographic coronary disease. Further, that when adding the subjects with previous CABG (some of whom may also represent recent, more acute conditions), the relation was even stronger.

In previous work [1], the aging of skin and vascular tissue has revealed a temporal concordance, including collagen abnormalities as well as incorporated lipid. Association between vascular disease and both skin cholesterol and ApoB levels has been observed [7]. While the pathophysiology of skin lipid remains to be fully explained [8], the opportunity to use skin bioassay as a surrogate vascular-disease marker appears viable and potentially important for screening algorithms, particularly if it corresponds to plaque stability.

These data suggest that while associated with the degree or presence of angiographic narrowing, elevated SkinT_c may also be associated with coronary-related events.

Table 1
Presentation characteristics by history of myocardial infarction (MI)

	Prior MI		<i>p</i> -Value
	No	Yes	
<i>N</i>	424	225	–
Age	62 ± 11	65 ± 12	0.001
Female	156 (37%)	58 (26%)	0.005
LDL-c (mg/dL)	118 ± 31	115 ± 33	0.24
HDL-c (mg/dL)	44 ± 14	38 ± 12	<0.001
History of hypertension	231 (54%)	133 (59%)	0.26
History of diabetes	58 (14%)	57 (25%)	<0.001
Smoker	125 (29%)	80 (36%)	0.11
Framingham score	8.3 ± 3.3	9.4 ± 3.8	<0.001
Number of vessels with >50% stenosis	0.9 ± 1.0	1.7 ± 1.0	<0.001
LAD > 50%	160 (38%)	143 (64%)	<0.001

Mean ± standard deviation for continuous measures; number and percent for categorical measures.

References

- [1] Lopukhin Y. *The skin and atherosclerosis (a three-drop test)*. Berkshire, UK: Harwood Academic Publishers; 1992. p. 3.
- [2] Sprecher DL, Goodman SG, Kannampuzha P, Pearce GL, Langer A. Skin tissue cholesterol (SkinT_c) is related to angiographically-defined cardiovascular disease. *Atherosclerosis* 2003;171:255–8.
- [3] Mancini GB, Chan S, Frohlich J, Kuramoto L, Schulzer M, Abbott D. Association of skin cholesterol content, measured by a noninvasive method, with markers of inflammation and Framingham risk prediction. *Am J Cardiol* 2002;89:1313–6.
- [4] Zawydiwski R, Sprecher DL, Eveleigh MJ, Horsewood P, Carte C, Patterson M. A novel test for the measurement of skin cholesterol. *Clin Chem* 2001;47:1302–4.
- [5] Wilson PW, D'Agostino RB, Levy D, Belanger A, Silbershatz H, Kannel W. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998;97:1837–47.
- [6] Ridker PM. Evaluating novel cardiovascular risk factors: can we better predict heart attacks? *Ann Intern Med* 1999;130:933–7.
- [7] Melico-Silvestre AA, Jacotot B, Buxtorf JC, Beaumont V, Beaumont JL. Study of free and esterified cholesterol in skin in atherogenic hyperlipidemias. *Pathol Biol (Paris)* 1981;29:573–8.
- [8] Tsuruoka H, Khovidhunkit W, Brown B, Fluhr JW, Elias PM, Feingold KR. Scavenger receptor class B type I is expressed in cultured keratinocytes and epidermis. *J Biol Chem* 2002;277:2916–22.